arXiv:math/0607094v2 [math.AT] 4 Apr 2007

SEMIFREE CIRCLE ACTIONS, BOTT TOWERS, AND QUASITORIC MANIFOLDS

MIKIYA MASUDA AND TARAS PANOV

Abstract. A *Bott tower* is the total space of a tower of fibre bundles with base $\mathbb{C}P^1$ and fibres $\mathbb{C}P^1$. Every Bott tower of height n is a smooth projective toric variety whose moment polytope is combinatorially equivalent to an n-cube. A circle action is *semifree* if it is free on the complement to fixed points. We show that a (quasi)toric manifold (in the sense of Davis–Januszkiewicz) over an n-cube with a semifree circle action and isolated fixed points is a Bott tower. Then we show that every Bott tower obtained in this way is topologically trivial, that is, homeomorphic to a product of 2-spheres. This extends a recent result of Ilinskii, who showed that a smooth compact toric variety with a semifree circle action and isolated fixed points is homeomorphic to a product of 2-spheres, and makes a further step towards our understanding of a problem motivated by Hattori's work on semifree circle actions. Finally, we show that if the cohomology ring of a quasitoric manifold (or Bott tower) is isomorphic to that of a product of 2-spheres, then the manifold is homeomorphic to the product.

1. Introduction

In their study of symmetric spaces Bott and Samelson [1] introduced a family of complex manifolds obtained as the total spaces of iterated bundles over $\mathbb{C}P^1$ with fibre $\mathbb{C}P^1$. Grossberg and Karshon [8] showed these manifolds to carry an algebraic torus action, therefore constituting an important family of smooth projective toric varieties, and called them *Bott towers*. The study of Bott towers has been significantly developed by Civan and Ray in [6] by enumerating the invariant stably complex structures and calculating their complex and real K-theory rings, and cobordisms.

An action of a group is called *semifree* if it is free in the complement to fixed points. A particularly interesting class of *Hamiltonian* semifree circle actions was studied by Hattori, who proved in [9] that a compact symplectic manifold M with a semifree Hamiltonian $\mathbb{S}^1$-action with non-empty isolated fixed point set has the same cohomology ring and the same Chern classes as $S^2\times\ldots\times S^2$, thus imposing a severe topological restriction on the topological structure of the manifold. Hattori's results were further extended by Tolman and Weitsman, who showed in [13] that a semifree symplectic $\mathbb{S}^1$-action with non-empty isolated fixed point set is automatically Hamiltonian, and the *equivariant* cohomology ring and Chern classes of M also agree with those of $S^2\times\ldots\times S^2$. In dimensions up to 6 it is known

---

1991 *Mathematics Subject Classification.* Primary 57S15; Secondary 14M25.

The second author gratefully acknowledges the support from the Japanese Society for the Promotion of Science (grant P05296), the Russian Foundation for Basic Research (grant 05-01-01032) and P. Deligne's 2004 Balzan prize in mathematics.

that a symplectic manifold with an $\mathbb{S}^1$-action satisfying the properties above is homeomorphic to a product of 2-spheres, but in higher dimensions the topological classification remains open.

Ilinskii considered in [16] an algebraic version of the above question on semifree symplectic $\mathbb{S}^1$-actions. Namely, he conjectured that a smooth compact complex algebraic variety X with a semifree action of the algebraic 1-torus $\mathbb{C}^*$ with positive number of isolated fixed points is homeomorphic to $\mathbb{C}P^1 \times \ldots \times \mathbb{C}P^1$. The algebraic and symplectic versions of the conjecture are related via the common subclass of projective varieties; a smooth projective variety is a symplectic manifold. Ilinskii proved the *toric* version of his algebraic conjecture, namely, the case when X is a *toric manifold* (a non-singular compact *toric variety*) and the semifree 1-torus is a subgroup of the acting torus (of dimension $\dim_{\mathbb{C}} X$). The first step of Ilinskii's argument was to show that if X admits a semifree subcircle with isolated fixed points, then the corresponding fan is combinatorially equivalent to the fan over the faces of a crosspolytope. A result of Dobrinskaya [15] implies that such X is a Bott tower, and this was the starting point in our study of semifree circle actions on Bott towers and related classes of manifolds with torus action, such as the *quasitoric manifolds*. This class of manifolds was introduced by Davis and Januszkiewicz in [7] (they used the name "toric manifolds", which we prefer to save for "non-singular compact toric varieties" as above). A quasitoric manifold is a compact manifold M of dimension $2n$ with a locally standard action of an n-dimensional torus $\mathbb{T}^n = \mathbb{S}^1 \times \ldots \times \mathbb{S}^1$ whose orbit space is a simple polytope P. Recently quasitoric manifolds have attracted substantial interest in the emerging field of "toric topology"; we review their construction in Section 3, for a more detailed account see [2, Ch. 5].

A projective toric manifold with the moment polytope P is a quasitoric manifold over P, and a Bott tower is a toric manifold with the moment polytope combinatorially equivalent to a cube (or simply a *toric manifold over a cube*). We therefore have the following hierarchy of classes of manifolds M with an action of $\mathbb{T}^n$:

$$\begin{aligned}\text{(1.1)}\quad \text{Bott towers} &\subset \text{Toric manifolds over cubes}\\ &\subset \text{Quasitoric manifolds over cubes.}\end{aligned}$$

By the above mentioned result of Dobrinskaya [15], the first inclusion above is in fact an identity (we explain this in Corollary 3.5). We proceed in Sections 4 and 5 by obtaining two results relating semifree circle actions on Bott towers, their topological structure, and cohomology rings. In Theorem 4.4 we show that if a Bott tower admits a semifree $\mathbb{S}^1$-action with isolated fixed points, then it is $\mathbb{S}^1$-equivariantly homeomorphic to a product of 2-spheres. We also show in Theorem 5.1 that a Bott tower with the cohomology ring isomorphic to that of a product of spheres is homeomorphic to the product. Both results are then extended to a much more general class of quasitoric manifolds over cubes (Theorems 4.8 and 5.7 respectively), which also allows us to deduce Ilinskii's result on semifree 1-dimensional algebraic torus actions on toric varieties (Corollary 4.9).

Since a cohomology isomorphism implies a homeomorphism in the case of product of spheres, we may ask a question of whether the cohomology ring detects the homeomorphism type of a Bott tower or quasitoric manifold in general. Some progress in this direction has been achieved in upcoming work [5] in the case of quasitoric manifolds over a product of simplices, which can be considered as an intermediate stage between quasitorics over a cube and the full generality.

It would be interesting to obtain smooth analogues of our classification results. In the case of Bott towers our homeomorphisms are actually diffeomorphisms (see Theorem 4.4 or Theorem 5.1), but some of our key arguments with quasitoric manifolds do not work in the smooth category. Although quasitoric manifolds are necessarily smooth [7, p. 421], the main problem here is that the original Davis and Januszkiewicz's classification result [7, Prop. 1.8] establishes only an equivariant *homeo*morphism between a quasitoric manifold and the canonical model determined by the polytope and the characteristic function. As the consequence, we don't know if there are exotic equivariant smooth structures, even on 4-dimensional quasitoric manifolds. (A canonical equivariant smooth structure, coinciding with the standard one in the toric case, is described in [3, §4].)

All the cohomology groups in the paper are taken with $\mathbb{Z}$ coefficients, unless otherwise specified.

The authors are grateful to Natalia Dobrinskaya and Dong Youp Suh for informal discussions of quasitoric manifolds over cubes and producs of simplices, and apologise for the improper references to [15] in the first version of the text. We also thank Takahiko Yoshida, who pointed out the difficulty in the smooth category for quasitoric manifolds. The second author wishes to thank Nigel Ray for introducing him to the study of Bott towers and illuminating discussions of the subject. Second author's thanks also go to Dmitri Timashev for drawing his attention to work of Ilinskii [16], which encouraged us to consider several related problems on semifree circle actions.

2. BOTT TOWERS

We briefly review the definitions here, following [8] and [6], where a reader may find a much more detailed account of the history and applications of Bott towers.

Definition. A *Bott tower* of height n is a sequence of manifolds $(B^{2k}\colon k \leqslant n)$ such that $B^2 = \mathbb{C}P^1$ and $B^{2k} = P(\underline{\mathbb{C}} \oplus \xi_{k-1})$ for $1 < k \leqslant n$ where $P(\cdot)$ denotes complex projectivisation, ξ_{k-1} is a complex line bundle over $B^{2(k-1)}$ and $\underline{\mathbb{C}}$ is a trivial line bundle. In particular, we have a bundle $B^{2k} \to B^{2(k-1)}$ with fibre $\mathbb{C}P^1$.

We shall also use the same name "Bott tower" for the last stage B^{2n} in the sequence; it follows from the definition that B^{2n} is a complex manifold obtained as the total space of an iterated bundle with fibre $\mathbb{C}P^1$. Bott towers of (real) dimension 4 are known as *Hirzebruch surfaces.*

The standard results on the cohomology of projectivised bundles lead to the following description of the cohomology ring of a Bott tower (see e.g. [6, Cor. 2.9]).

Lemma 2.1. *The cohomology of B^{2k} is a free module over $H^*(B^{2(k-1)})$ on generators 1 and u_k, which have dimensions 0 and 2 respectively; the ring* *structure is determined by the single relation*

$$u_k^2 = c_1(\xi_{k-1})u_k,$$

and the restriction of u_k *to the fibre* $\mathbb{C}P^1 \subset B^{2k}$ *is the first Chern class of the canonical line bundle over* $\mathbb{C}P^1$*.*

We set u_1 to be the canonical generator of $H^2(\mathbb{C}P^1)$ (the first Chern class of the canonical line bundle). The Bott tower is therefore determined by the list of integers $(a_{ij}\colon 1 \leqslant i < j \leqslant n)$, where

$$u_k^2 = \sum_{i=1}^{k-1} a_{ik}u_iu_k, \qquad 1 \leqslant k \leqslant n. \tag{2.1}$$

The cohomology ring of B^{2n} is the quotient of $\mathbb{Z}[u_1, \dots, u_n]$ by relations (2.1).

It is convenient to organise the integers a_{ij} into an $n \times n$ upper triangular matrix,

$$A = \begin{pmatrix} -1 & a_{12} & \cdots & a_{1n} \\ 0 & -1 & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -1 \end{pmatrix}. \tag{2.2}$$

Example 2.2. When $n = 2$, the Bott tower B^4 is determined by a single line bundle ξ_1 over $\mathbb{C}P^1$. We have $\xi_1 = \gamma^m$ for some $m \in \mathbb{Z}$ where γ is the canonical line bundle over $B^2 = \mathbb{C}P^1$, and so the cohomology ring is determined by the relations $u_1^2 = 0$ and $u_2^2 = mu_1u_2$. It is well-known that

$$P(\underline{\mathbb{C}} \oplus \gamma^m) \cong P(\underline{\mathbb{C}} \oplus \gamma^{m'}) \text{ if and only if } m \equiv m' \pmod 2,$$

where $\cong$ means "diffeomorphic". The proof goes as follows. We note that $P(E) \cong P(E \otimes \eta)$ for any complex line bundle η. Suppose $m \equiv m' \pmod 2$. Then $m' - m = 2\ell$ for some $\ell \in \mathbb{Z}$ and we have diffeomorphisms

$$P(\underline{\mathbb{C}} \oplus \gamma^m) \cong P((\underline{\mathbb{C}} \oplus \gamma^m) \otimes \gamma^\ell) = P(\gamma^\ell \oplus \gamma^{m+\ell}).$$

Here both $\gamma^\ell \oplus \gamma^{m+\ell}$ and $\underline{\mathbb{C}} \oplus \gamma^{m'}$ are over $\mathbb{C}P^1$ and have equal first Chern class, so they are isomorphic. Hence the last space above is $P(\underline{\mathbb{C}} \oplus \gamma^{m'})$. On the other hand, it is not difficult to see that if $H^*(P(\underline{\mathbb{C}} \oplus \gamma^m)) \cong H^*(P(\underline{\mathbb{C}} \oplus \gamma^{m'}))$ as rings, then $m \equiv m' \pmod 2$.

This example shows that the cohomology ring determines the topological type of a Bott tower B^{2n} for $n = 2$. A case-to-case analysis based on a classification result of [15, §3] shows that this is also the case for $n = 3$. So we may ask the following question.

Problem 2.3. *Are Bott towers* B_1^{2n} *and* B_2^{2n} *homeomorphic if* $H^*(B_1^{2n}) \cong H^*(B_2^{2n})$ *as rings?*

We investigate this question further in Section 5, where a partial answer is given.

3. Quasitoric manifolds

Davis and Januszkiewicz introduced in [7] a class of $2n$-dimensional manifolds M with an action of n-dimensional torus T. They required the action to be *locally standard* (locally isomorphic to the standard T-representation in $\mathbb{C}^n$) and the quotient space M/T to be homeomorphic to a *simple* n-dimensional polytope P, so there is a projection $\pi\colon M \to P$ whose fibres are orbits of the action. Davis and Januszkiewicz describe their manifolds as *toric*; more recently, the term *quasitoric* has been adopted, to avoid confusion with the non-singular compact toric varieties of algebraic geometry. We follow this convention below, and refer to such M as *quasitoric manifolds*, saving the term "toric manifolds" to describe non-singular compact toric varieties. We note that a projective toric manifold is a quasitoric manifold; for more discussion of the relationship between the two classes see [2, Ch. 5].

Let m denote the number of *facets* (codimension-one faces) of P; we order the facets so that the first n of them meet at a vertex. We denote the facets by F_i for $1 \leqslant i \leqslant m$, and denote $\mathcal{F}$ the set of all facets. The preimage $\pi^{-1}(F_i)$ is a connected codimension-two submanifold of M, fixed pointwise by a circle subgroup of T. We denote it by M_i and refer to it as the *characteristic submanifold* corresponding to F_i, for $1 \leqslant i \leqslant m$. An *omniorientation* [4] of M consists of a choice of orientation for M and for each characteristic submanifold.

Let N denote the integer lattice of one-parameter circle subgroups in T, so we have $N \cong \mathbb{Z}^n$. Given a characteristic submanifold M_j, we denote by λ_j a primitive vector in N that spans the circle subgroup $T_{M_j} \subset T$ fixing M_j. The vector λ_j is determined up to sign. The correspondence $\lambda\colon F_j \mapsto \lambda_j$ was called in [7] the *characteristic function* corresponding to M.

The omniorientation allows us to interpret the characteristic function as a linear map $\lambda\colon \mathbb{Z}^{\mathcal{F}} \to N$. This is done as follows. First, we notice that an action of a one-parameter circle subgroup of T determines an orientation for the normal bundle ν_j of the embedding $M_j \subset M$. The omniorientation of M also provides an orientation for ν_j by means of the following decomposition of the tangent bundle:

$$\tau(M)|_{M_j} = \tau(M_j) \oplus \nu_j.$$

Now we choose the primitive vectors λ_j so that the two orientations of ν_j coincide for $1 \leqslant j \leqslant m$.

In general, there is no canonical choice of an omniorientation for M. However, if M admits a *T-equivariant almost complex structure*, then a choice of such a structure provides a canonical way of orienting M and normal bundles ν_j for $1 \leqslant j \leqslant m$, thereby specifying an omniorientation *associated* with the equivariant almost complex structure. In what follows we shall always choose the associated omniorientation if M is equivariantly almost complex (in particular, if M is a toric manifold); otherwise we shall fix an arbitrary omniorientation.

By definition, the characteristic function satisfies the *non-singularity condition*: $\lambda_{j_1}, \dots, \lambda_{j_n}$ is a basis of N whenever the intersection $F_{j_1} \cap \dots \cap F_{j_n}$ is non-empty. So we may use the vectors $\lambda_1, \dots, \lambda_n$ to define a basis for N, thereby identifying it with $\mathbb{Z}^n$, and represent the map λ by an $n \times m$ integral matrix of the form

$$\Lambda = \begin{pmatrix} 1 & 0 & \dots & 0 & \lambda_{1,n+1} & \dots & \lambda_{1,m} \\ 0 & 1 & \dots & 0 & \lambda_{2,n+1} & \dots & \lambda_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & \lambda_{n,n+1} & \dots & \lambda_{n,m} \end{pmatrix}. \tag{3.1}$$

It is often convenient to partition Λ as $(E \mid \Lambda_\star)$, where E is a unit matrix and $\Lambda_\star$ is $n \times (m-n)$. For any other vertex $F_{j_1} \cap \dots \cap F_{j_n}$ the corresponding columns $\lambda_{j_1}, \dots, \lambda_{j_n}$ form a basis for $\mathbb{Z}^n$, and have determinant ± 1. We refer to (3.1) as the *refined form* of characteristic matrix Λ, and call $\Lambda_\star$ its *reduced submatrix.*

Having chosen a basis for N, we may identify our torus T with the standard product of unit circles in $\mathbb{C}^n$:

$$\mathbb{T}^n = \{(e^{2\pi i \varphi_1}, \dots, e^{2\pi i \varphi_n}) \in \mathbb{C}^n\}, \tag{3.2}$$

where $(\varphi_1, \dots, \varphi_n)$ runs over $\mathbb{R}^n$. We shall also denote a generic point of $\mathbb{T}^n$ by $(t_1, \dots, t_n)$. The circle subgroup fixing M_j can now be written as

$$T_{M_j} = \{(t^{\lambda_{1j}}, \dots, t^{\lambda_{nj}}) = (e^{2\pi i \lambda_{1j}\varphi}, \dots, e^{2\pi i \lambda_{nj}\varphi}) \in \mathbb{T}^n\}, \quad 1 \leqslant j \leqslant m \tag{3.3}$$

where $\varphi \in \mathbb{R}$ and $t = e^{2\pi i \varphi}$.

Remark. Not every toric manifold X is a quasitoric manifold, as the quotient X/T may fail to be a simple polytope (although it is a simple polytope when X is projective). Nevertheless, X has characteristic submanifolds X_j (T-invariant divisors), and there is a canonical omniorientation induced from the complex structures on X and X_j. Therefore, the characteristic matrix Λ and its reduced submatrix $\Lambda_\star$ are defined for every toric manifold X. The vectors λ_j are the primitive vectors along the edges of the fan corresponding to X.

Let v_j denote the class in $H^2(M)$ dual to the fundamental class of M_j, $1 \leqslant j \leqslant m$. According to [7, Th. 4.14], the ring $H^*(M)$ is generated by $v_1, \dots, v_m$, modulo two sets of relations. The first set is formed by the monomial relations which arise from the *Stanley–Reisner ideal* of P; the second set consists of linear relations determined by the characteristic matrix:

$$v_i = -\lambda_{i,n+1}v_{n+1} - \dots - \lambda_{i,m}v_m \quad \text{for} \quad 1 \leqslant i \leqslant n. \tag{3.4}$$

It follows that $v_{n+1}, \dots, v_m$ suffice to generate $H^*(M)$ multiplicatively.

Two quasitoric manifolds M_1 and M_2 are said to be *weakly T-equivariantly homeomorphic* (or simply *weakly T-homeomorphic*) if there is an automorphism $\theta\colon T \to T$ and a homeomorphism $f\colon M_1 \to M_2$ such that $f(t \cdot x) = \theta(t) \cdot f(x)$ for every $t \in T$ and $x \in M_1$. If θ is identity, then M_1 and M_2 are *T-homeomorphic*. Following Davis and Januszkiewicz, we say that two quasitoric manifolds M_1 and M_2 over the same P are *equivalent* if there is a weak T-homeomorphism $f\colon M_1 \to M_2$ covering the identity on P. By [7, Prop. 1.8], a quasitoric manifold M over P is determined up to equivalence by its characteristic function λ. This follows from the "basic construction" providing a quasitoric manifold $M(\lambda)$, which depends only on P and λ, together with a T-equivariant homeomorphism $M(\Lambda) \to M$ covering the identity on P.

Let $\mathrm{cf}(P)$ denote the set of characteristic functions on the facets of P, that is, the set of maps $\lambda\colon \mathcal{F} \to N$ satisfying the non-singularity condition. The group $GL(N) \cong GL(n, \mathbb{Z})$ of automorphisms of the lattice N acts on the set $\mathrm{cf}(P)$ from the left (an automorphism $g\colon N \to N$ acts by composition $\lambda \mapsto g \cdot \lambda$). Since automorphisms of the lattice N correspond to automorphisms of the torus T, there is a one-to-one correspondence

$$GL(N)\backslash \mathrm{cf}(P) \longleftrightarrow \{\text{equivalence classes of } M \text{ over } P\}. \tag{3.5}$$

One may assign an $n \times m$-matrix Λ to an element $\lambda \in \mathrm{cf}(P)$ by ordering the facets and choosing a basis for N, as we did above. A choice of matrix $\mathit{\Lambda}$ in the refined form (3.1) can now be regarded as a choice of a specific representative of the left coset class in $GL(N)\backslash \mathrm{cf}(P)$. If a characteristic matrix is given in an unrefined form $\mathit{\Lambda} = (A \mid B)$, where A is $n \times n$ and B is $n \times (m - n)$, then the refined representative in its coset class is given by $(E \mid A^{-1}B)$.

A quasitoric manifold over P can be obtained as the quotient of the *moment-angle manifold* $\mathcal{Z}_P$ by a freely acting $(m-n)$-dimensional torus subgroup of $\mathbb{T}^m$ determined by the kernel of the characteristic map $\lambda\colon \mathbb{Z}^m \to N$, see [7], [2, Ch. 6]. The moment-angle manifold $\mathcal{Z}_P$ embeds into $\mathbb{C}^m$ as a complete intersection of $m-n$ real quadratic hypersurfaces [3, §3]. It follows that both $\mathcal{Z}_P$ and M are necessarily smooth, but we don't know if this equivariant smooth structure on M is unique (see the discussion in the Introduction).

We are particularly interested in the case when the quotient polytope $P = M/\mathbb{T}^n$ is the n-cube $\mathbb{I}^n$. Then $m = 2n$ and we shall additionally assume that the facets F_j and F_{n+j} are opposite (i.e., do not intersect) for $1 \leqslant j \leqslant n$. In the case $P = \mathbb{I}^n$ the moment-angle manifold is the product of n three-dimensional spheres, embedded in $\mathbb{C}^{2n}$ as

$$\{(z_1, \dots, z_{2n}) \in \mathbb{C}^{2n}\colon |z_j|^2 + |z_{n+j}|^2 = 1 \text{ for } 1 \leqslant j \leqslant n\}.$$

The quotient $(S^3)^n/\mathbb{T}^{2n}$ by the coordinatewise action is a cube $\mathbb{I}^n$. The n-dimensional subtorus $T(\mathit{\Lambda}) \subset \mathbb{T}^{2n}$ determined by the kernel of characteristic map (3.1) is given by

$$\begin{aligned}(t_1, \dots, t_n) \mapsto (&t_1^{-\lambda_{1,n+1}} t_2^{-\lambda_{1,n+2}} \cdots t_n^{-\lambda_{1,2n}}, \dots, \\ &t_1^{-\lambda_{n,n+1}} t_2^{-\lambda_{n,n+2}} \cdots t_n^{-\lambda_{n,2n}}, t_1, t_2, \dots, t_n).\end{aligned} \tag{3.6}$$

It acts freely on $(S^3)^n$, and the quotient $(S^3)^n/T(\mathit{\Lambda})$ is the quasitoric manifold M determined by $\mathit{\Lambda}$. The n-dimensional torus $\mathbb{T}^{2n}/T(\mathit{\Lambda}) \cong \mathbb{T}^n$ acts on $(S^3)^n/T(\mathit{\Lambda}) \cong M$ with quotient $\mathbb{I}^n$. In coordinates, $(t_1, \dots, t_n) \in \mathbb{T}^n$ acts on the equivalence class $[z_1, \dots, z_{2n}] \in (S^3)^n/T(\mathit{\Lambda})$ as multiplication by $(t_1, \dots, t_n, 1, \dots, 1)$.

Proposition 3.1. *A Bott tower carries a natural torus action turning it into a quasitoric manifold over a cube with the reduced submatrix* $\mathit{\Lambda}_\star = A^t$*, see* (2.2) *and* (3.1).

Proof. As it is shown in [6, Prop. 3.1], the Bott tower corresponding to (2.2) can be obtained as the quotient of $(S^3)^n$ by an n-dimensional subtorus of $\mathbb{T}^{2n}$ defined by the inclusion

$$(t_1, \dots, t_n) \mapsto (t_1, t_1^{-a_{12}} t_2, \dots, t_1^{-a_{1n}} t_2^{-a_{2n}} \cdots t_{n-1}^{-a_{n-1,n}} t_n, t_1, t_2, \dots, t_n).$$

It remains to note that this coincides with $T(\Lambda)$ from (3.6) for $\Lambda_\star = A^t$. $\square$

Remark. The Stanley–Reisner relations for the n-cube are $v_i v_{i+n} = 0$, $1 \leqslant i \leqslant n$. These relations together with (3.4) give (2.1) for $\Lambda_\star = A^t$ and $u_i = v_{i+n}$, $1 \leqslant i \leqslant n$.

By definition, a Bott tower is a complex manifold. A non-compact version of (3.6) was used in [8] to describe Bott towers as non-singular projective toric varieties. The two approaches are related in [6, §2].

Given a permutation σ of n elements, denote by $P(\sigma)$ the corresponding $n \times n$ *permutation matrix*, which has units in positions $(\sigma(i), i)$ for $1 \leqslant i \leqslant n$, and zeros elsewhere. There is an action of the symmetric group on $n \times n$ matrices by conjugations $A \mapsto P(\sigma)^{-1} A P(\sigma)$ or, equivalently, by permutations of the rows and columns of A.

Proposition 3.2. *A quasitoric manifold M over a cube with reduced submatrix $\Lambda_\star$ is equivalent to a Bott tower if and only if $\Lambda_\star$ can be conjugated by a permutation matrix to an upper triangular matrix.*

Proof. Assume that $\Lambda_\star$ can be conjugated by a permutation matrix to an upper triangular matrix. Consider the action of S_n on the set of facets of $\mathbb{I}^n$ by permuting the pairs of opposite facets. A reordering of facets corresponds to reordering of columns in the $n \times 2n$ characteristic matrix Λ, so an element $\sigma \in S_n$ acts as

$$\Lambda \mapsto \Lambda \cdot \begin{pmatrix} P(\sigma) & 0 \\ 0 & P(\sigma) \end{pmatrix}.$$

This action does not preserve the refined form of Λ, as $(E \mid \Lambda_\star)$ becomes $(P(\sigma) \mid \Lambda_\star P(\sigma))$. The refined representative in the left coset class (3.5) of the latter matrix is given by $(E \mid P(\sigma)^{-1} \Lambda_\star P(\sigma))$. (In other words, we have to compensate the permutation of pairs of facets by an automorphism of $\mathbb{T}^n$ permuting the coordinate subcircles to keep the characteristic matrix in the refined form.) This implies that the permutation action on the pairs of opposite facets induces the conjugation action on the reduced submatrices. So we may assume, up to equivalence, that M has an upper triangular reduced submatrix $\Lambda_\star$. The non-singularity condition guarantees that the diagonal entries of $\Lambda_\star$ are ± 1, and we can set all of them to -1 by changing the omniorientation of M if necessary. Now, M and the Bott tower corresponding to the matrix $A = \Lambda_\star^t$ have the same characteristic matrices Λ by Proposition 3.1. Therefore, they are equivalent by [7, Prop. 1.8]. The opposite statement follows from Proposition 3.1. $\square$

It is now clear that not all quasitoric manifolds over a cube are Bott towers. For example, a 4-dimensional quasitoric manifold over a square with reduced submatrix $\Lambda_\star = \begin{pmatrix} -1 & -2 \\ -1 & -1 \end{pmatrix}$ is not a Bott tower, as $\Lambda_\star$ cannot be conjugated to an upper triangular matrix. (The corresponding manifold is homeomorphic to $\mathbb{C}P^2 \# \mathbb{C}P^2$, and therefore does not even admit a complex structure [7].)

Given a k-element subset $\{i_1, \ldots, i_k\}$ of n elements, the corresponding *principal minor* of a square n-matrix A is the determinant of the submatrix formed by elements in columns and rows $i_1, \ldots, i_k$. For Bott towers, Proposition 3.1 ensures that all principal minors of the matrix $-\Lambda_\star$ equal 1; while for arbitrary quasitoric manifolds the non-singularity condition only guarantees that every principal minor of $\Lambda_\star$ is ± 1.

Recall that an upper triangular matrix is *unipotent* if all its diagonal entries are unit. The following key technical lemma can be retrieved from the proof of a much more general result of Dobrinskaya [15, Th. 6]. We give a slightly more expanded proof here for the sake of completeness.

Lemma 3.3 ([15]). *Let R be a commutative integral domain with an identity element 1, and let A be an $n \times n$ matrix with entries in R. Suppose that every proper principal minor of A is 1. If $\det A = 1$, then A is conjugate by a permutation matrix to a unipotent upper triangular matrix, and otherwise to a matrix of the form*

$$\begin{pmatrix} 1 & b_1 & 0 & \dots & 0 \\ 0 & 1 & b_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1} \\ b_n & 0 & \dots & 0 & 1 \end{pmatrix} \tag{3.7}$$

where $b_i \neq 0$ for every i.

Proof. Every diagonal entry of A must be 1. We say that the i-th row is *elementary* if its i-th entry is 1 and others are 0. Assuming by induction that the theorem holds for matrices of size $(n-1)$, we deduce that A itself is conjugate to a unipotent upper triangular matrix if and only if it contains an elementary row. Denote by A_i the square matrix of size $(n-1)$ obtained by removing the i-th column and row from A.

We may assume by induction that A_n is a unipotent upper triangular matrix. Now we apply the induction assumption to A_1. The permutation of rows and columns turning A_1 into a unipotent upper triangular matrix turns A into an "almost" unipotent upper triangular matrix; the latter may have only one non-zero entry below the diagonal, and it must be in the first column. If this non-zero entry is not a_{n1}, then the n-th row of A is elementary, and A is conjugate to a unipotent upper triangular matrix. Therefore, we may assume that

$$A = \begin{pmatrix} 1 & * & * & \dots & * \\ 0 & 1 & * & \dots & * \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1} \\ b_n & 0 & \dots & 0 & 1 \end{pmatrix},$$

where $b_{n-1} \neq 0$ and $b_n \neq 0$ (otherwise A has an elementary row). Now let a_{1j_1} be the last non-zero non-diagonal entry in the first row of A. If A does not have an elementary row, the we may define inductively $a_{j_i j_{i+1}}$ as the last non-zero non-diagonal entry in the j_i-th row of A. Clearly, we have

$$1 < j_1 < \ldots < j_i < j_{i+1} < \ldots < j_k = n$$

for some $k < n$. Now, if $j_i = i + 1$ for $i = 1, \dots, n-1$, then A is matrix (3.7) with $b_i = a_{j_{i-1}j_i}$, $i = 1, \dots, n-1$. Otherwise, the submatrix

$$S = \begin{pmatrix} 1 & a_{1j_1} & 0 & \dots & 0 \\ 0 & 1 & a_{j_1j_2} & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & a_{j_{k-1}n} \\ b_n & 0 & \dots & 0 & 1 \end{pmatrix}$$

of A, formed by columns and rows $1, j_1, \dots, j_k$, is proper and has determinant $1 \pm b_n \prod a_{j_ij_{i+1}} \neq 1$. This contradiction finishes the proof. □

The following theorem is not new; the equivalence *a*) ⇔ *b*) is a particular case of [15, Th. 6], while *b*) ⇔ *c*) was discussed (in a more general case of quasitoric manifolds over arbitrary polytopes) in [11] and [2, Prop. 5.53]. We give a proof because it is needed in the next sections.

Theorem 3.4. *Let M be a quasitoric manifold over a cube, and $\Lambda_\star$ the corresponding reduced submatrix. The following conditions are equivalent:*

a) *M is equivalent to a Bott tower;*
b) *all principal minors of $-\Lambda_\star$ are 1;*
c) *M has a $\mathbb{T}^n$-equivariant almost complex structure (with the associated omniorientation).*

Proof. The implication b) ⇒ a) follows from Lemma 3.3 and Proposition 3.2. The implication a) ⇒ c) is obvious. Let us prove c) ⇒ b). First, we recall from [11, §4], [15] and [17] the definition of the *sign* $\sigma(p)$ for a fixed point of the $\mathbb{T}^n$-action on M. Every fixed point p can be obtained as the intersection $M_{j_1} \cap \ldots \cap M_{j_n}$ of n characteristic submanifolds, and corresponds to the vertex of P obtained as the intersection $F_{j_1} \cap \ldots \cap F_{j_n}$ of the corresponding facets. The tangent space to M at p therefore decomposes into the sum of normal subspaces to M_{j_k} for $1 \leqslant k \leqslant n$:

$$\tau_p(M) = \nu_{j_1}\big|_p \oplus \ldots \oplus \nu_{j_n}\big|_p. \tag{3.8}$$

The omniorientation of M provides two different ways of orienting the above space; we set $\sigma(p) = 1$ if these two orientations coincide and $\sigma(p) = -1$ otherwise. This sign can be calculated in terms of P and characteristic matrix Λ as

$$\sigma(p) = \operatorname{sign}\bigl(\det(\lambda_{j_1}, \ldots, \lambda_{j_n}) \cdot \det(a_{j_1}, \ldots, a_{j_n})\bigr) \tag{3.9}$$

(see [17, §1]), where a_i denotes a normal vector to the facet F_i pointing inside the polytope. Now, if $P = \mathbb{I}^n$, then every fixed point p corresponds to a vertex given as

$$F_{i_1} \cap \ldots \cap F_{i_k} \cap F_{n+l_1} \cap \ldots \cap F_{n+l_{n-k}}$$

for some $1 \leqslant i_1 < \ldots < i_k \leqslant n$ and $1 \leqslant l_1 < \ldots < l_{n-k} \leqslant n$, and we may choose $a_i = e_i$ (the ith basis vector) for $1 \leqslant i \leqslant n$ and $a_j = -e_j$ for $n+1 \leqslant j \leqslant 2n$. Thus, the expression in the right hand side of (3.9) equals the principal minor of $-\Lambda_\star$ formed by the columns and rows with numbers $l_1, \ldots, l_{n-k}$. It remains to notice that in the almost complex case the two orientations in (3.8) coincide, so the sign of every fixed point is 1. □

Recall that a *crosspolytope* is a regular polytope dual to the cube (in particular, a 3-dimensional crosspolytope is an octahedron).

Corollary 3.5. *Let X be a toric manifold whose associated fan is combinatorially equivalent to the fan consisting of the cones over the faces of a crosspolytope. Then X is a Bott tower.*

Proof. The reduced matrix $\Lambda_\star$ of X is $n \times n$ and all principal minors of $-\Lambda_\star$ are unit by the same reason as in the proof of Theorem 3.4. By Lemma 3.3, $-\Lambda_\star$ can be conjugated to a unipotent uppper triangular matrix, so Λ is of the same form as the characteristic matrix of a Bott tower. In the toric manifold case, the columns of Λ are the primitive vectors along the edges of the fan, so the combinatorial type of the fan and Λ completely determine the fan. It follows that the fan of X is the same as the fan of a Bott tower, which implies that X is a Bott tower by the one-to-one correspondence between fans and toric varieties. □

A toric manifold over a cube satisfies the assumption of Corollary 3.5. It follows that the class of Bott towers coincides with the class of toric manifolds over a cube, and the first inclusion in (1.1) is an identity. Like Lemma 3.3, Corollary 3.5 is a particular case of a more general result of Dobrinskaya [15, Cor. 7], which gives a criterion for a quasitoric manifold over a product of simplices to be decomposable into a tower of fibre bundles.

4. Semifree circle actions

An action of a group on a topological space is called *semifree* if it is free on the complement to fixed points. We first show (Theorem 4.3 below) that if the torus $\mathbb{T}^n$ acting on a quasitoric manifold M over a cube has a circle subgroup acting semifreely and with isolated fixed points, then M is a Bott tower. Then we prove that all these Bott towers are $\mathbb{S}^1$-equivariantly homeomorphic to a product of 2-dimensional spheres (with the diagonal $\mathbb{S}^1$-action).

A complex n-dimensional representation of $\mathbb{S}^1$ is determined by a set of weights $k_j \in \mathbb{Z}$ for $1 \leqslant j \leqslant n$. In appropriate coordinates, an element $s = e^{2\pi i\varphi} \in \mathbb{S}^1$ acts as

$$s \cdot (z_1, \ldots, z_n) = (e^{2\pi i k_1\varphi} z_1, \ldots, e^{2\pi i k_n\varphi} z_n). \tag{4.1}$$

The following statement is straightforward.

Proposition 4.1. *A representation of $\mathbb{S}^1$ in $\mathbb{C}^n$ is semifree if and only if $k_j = \pm 1$ for $1 \leqslant j \leqslant n$.*

A circle subgroup in $\mathbb{T}^n$ is determined by an integer primitive vector $\nu = (\nu_1, \ldots, \nu_n)$:

$$S^1_\nu = \{(e^{2\pi i\nu_1\varphi}, \ldots, e^{2\pi i\nu_n\varphi})\} \subseteq \mathbb{T}^n. \tag{4.2}$$

We shall consider the tangential representations of $\mathbb{T}^n$ and its circle subgroups at fixed points of M. The representation of $\mathbb{T}^n$ in the tangent space $\tau_p(M)$ at a fixed point $p = M_{j_1} \cap \ldots \cap M_{j_n}$ decomposes into the sum of non-trivial real two-dimensional representations in the normal subspaces ν_{j_k} of M_{j_k}. The omniorientation endows each ν_{j_k} with a complex structure, therefore identifying it with $\mathbb{C}$ and τ_pM with $\mathbb{C}^n$. In these coordinates, the weights of the representation of circle subgroup (4.2) in τ_pM can be identified with the coefficients $k_i = k_i(\nu, p)$, $1 \leqslant i \leqslant n$, of the decomposition of ν in terms of $\lambda_{j_1}, \ldots, \lambda_{j_n}$:

$$\nu = k_1(\nu,p)\lambda_{j_1} + \ldots + k_n(\nu,p)\lambda_{j_n} \tag{4.3}$$

(see, e.g. [17, Lemma 2.3]).

Corollary 4.2. *A subcircle* $S^1_\nu \subseteq \mathbb{T}^n$ *acts on a quasitoric manifold* M *semifreely and with isolated fixed points if and only if for every vertex* $p = F_{j_1} \cap \ldots \cap F_{j_n}$ *the coefficients in* (4.3) *satisfy* $k_i(\nu, p) = \pm 1$ *for* $1 \leqslant i \leqslant n$.

Theorem 4.3. *Let* M *be a quasitoric manifold over a cube with reduced submatrix* $\varLambda_\star$. *Assume that* M *admits a semifree circle subgroup with isolated fixed points. Then* M *is equivalent to a Bott tower.*

Proof. We may assume by induction that every characteristic submanifold is a Bott tower, so the determinant of every proper principal minor of $-\varLambda_\star$ is 1. Therefore, we are in the situation of Lemma 3.3, and $-\varLambda_\star$ is of one of the two types described there. The second type is ruled out because of the semifree assumption. Indeed, assume that $\varLambda = (E \mid B)$ where B is matrix (3.7), and $S^1_\nu \subseteq \mathbb{T}^n$ acts semifreely with isolated fixed points. Applying the criterion from Corollary 4.2 to the vertex $p = F_1 \cap \ldots \cap F_n$, we obtain $\nu_i = \pm 1$ for $1 \leqslant i \leqslant n$. Now apply the same criterion to the vertex $p' = F_{n+1} \cap \ldots \cap F_{2n}$. Since the submatrix formed by the corresponding columns of $\varLambda$ is exactly B, we have $\det B = \pm 1$. This implies that at least one of b_i equals ± 1, i.e. at least one of the rows of B consists only of two ± 1's and zeros. Therefore, if all the coefficients $k_i(\nu, p')$ in the expression $\nu = k_1(\nu,p')\lambda_{n+1} + \ldots + k_n(\nu,p')\lambda_{2n}$ are ± 1, then at least one component ν_i in the left hand side is even. A contradiction. □

Our next result shows that a Bott tower with a semifree circle subgroup and isolated fixed points is topologically (or even $\mathbb{S}^1$-equivariantly) trivial, i.e. homeomorphic to a product of 2-spheres. Let t (respectively, $\mathbb{C}$) denote the standard (respectively, trivial) complex one-dimensional $\mathbb{S}^1$-representation. A product bundle with fibre V will be denoted by $\underline{V}$. We say that an action of a group G on a Bott tower B^{2n} *preserves the tower structure* if for each stage $B^{2k} = P(\underline{\mathbb{C}} \oplus \xi_{k-1})$, $k \leqslant n$, the line bundle ξ_{k-1} is G-equivariant. The intrinsic $\mathbb{T}^n$-action obviously preserves the tower structure.

Theorem 4.4. *Assume that a Bott tower* B^{2n} *admits a semifree* $\mathbb{S}^1$*-action with isolated fixed points preserving the tower structure. Then* B^{2n} *is* $\mathbb{S}^1$*-equivariantly diffeomorphic to the product* $(P(\mathbb{C} \oplus t))^n$.

Proof. We may assume by induction that the $(n-1)$-stage of the Bott tower is $(P(\mathbb{C} \oplus t))^{n-1}$ and $B^{2n} = P(\underline{\mathbb{C}} \oplus \xi)$ for some $\mathbb{S}^1$-line bundle ξ over $(P(\mathbb{C} \oplus t))^{n-1}$.

Let γ be the unique $\mathbb{S}^1$-line bundle over $P(\mathbb{C} \oplus t) \cong \mathbb{C}P^1$ whose underlying bundle is the canonical line bundle and

$$\gamma|_{(1:0)} = \mathbb{C} \qquad \text{and} \qquad \gamma|_{(0:1)} = t. \tag{4.4}$$

Denote by $x \in H^2(P(\mathbb{C} \oplus t))$ the first Chern class of γ, and let $x_i \in H^2(P(\mathbb{C} \oplus t)^{n-1})$ be the pullback of x by the projection π_i onto the i-th factor. The first Chern class of ξ may then be written as $\sum_{i=1}^{n-1} a_i x_i$ with $a_i \in \mathbb{Z}$. The $\mathbb{S}^1$-line bundles ξ and $\otimes_{i=1}^{n-1}\pi_i^*(\gamma^{a_i})$ have the same underlying bundles; so there is an integer k such that

$$\xi = t^k \bigotimes_{i=1}^{n-1} \pi_i^*(\gamma^{a_i}) \tag{4.5}$$

as $\mathbb{S}^1$-line bundles, see [10, Cor. 4.2].

We encode the fixed points of $P(\mathbb{C}\oplus t)^{n-1}$ as $(p_1^{\epsilon_1}, \dots, p_{n-1}^{\epsilon_{n-1}})$ where $\epsilon_i = 0$ or 1 and $p_i^{\epsilon_i}$ denotes $(1:0)$ if $\epsilon_i = 0$ and $(0:1)$ if $\epsilon_i = 1$. It follows from (4.4) and (4.5) that

$$\xi|_{(p_1^{\epsilon_1}, \dots, p_{n-1}^{\epsilon_{n-1}})} = t^{k+\sum_{i=1}^{n-1}\epsilon_i a_i}.$$

The $\mathbb{S}^1$-action on $B^{2n} = P(\underline{\mathbb{C}}\oplus\xi)$ is semifree if and only if $|k+\sum_{i=1}^{n-1}\epsilon_i a_i| = 1$ for all possible values of ϵ_i's. Setting $\epsilon_i = 0$ for all i we get $|k| = 1$. Assume $k = 1$ (the case $k = -1$ is treated similarly). Then $(a_1, \dots, a_{n-1}) = (0, \dots, 0)$ or $(0, \dots, 0, -2, 0, \dots, 0)$. In the former case, $\xi = \underline{t}$ and $B^{2n} = P(\underline{\mathbb{C}}\oplus\xi) \cong P(\mathbb{C}\oplus t)^n$. In the latter case we have $\xi = t\pi_i^*(\gamma^{-2})$ for some i, so that $B^{2n} = P(\underline{\mathbb{C}}\oplus\xi)$ is the pullback of $P(\underline{\mathbb{C}}\oplus t\gamma^{-2})$ by the projection π_i. Since for any $\mathbb{S}^1$-vector bundle E and $\mathbb{S}^1$-line bundle η the projectivisations $P(E)$ and $P(E\otimes\eta)$ are $\mathbb{S}^1$-diffeomorphic, we have $P(\underline{\mathbb{C}}\oplus t\gamma^{-2}) \cong P(\gamma\oplus t\gamma^{-1})$. The first Chern class of $\gamma\oplus t\gamma^{-1}$ is zero, so its underlying bundle is trivial. The $\mathbb{S}^1$-representation in the fibre of $\gamma\oplus t\gamma^{-1}$ over a fixed point is $\mathbb{C}\oplus t$ by (4.4). Therefore, $\gamma\oplus t\gamma^{-1} = \underline{\mathbb{C}}\oplus\underline{t}$ as $\mathbb{S}^1$-bundles. It follows that $P(\underline{\mathbb{C}}\oplus t\gamma^{-2}) \cong P(\underline{\mathbb{C}}\oplus\underline{t})$, which finishes the proof. ☐

Remark. The diffeomorphism of Theorem 4.4 is not a diffeomorphism of $\mathbb{T}^n$-manifolds.

Our next aim is to generalise Theorem 4.4 to quasitoric manifolds. Although the result does not hold for all quasitoric manifolds (see the next Example), it remains true if we additionally require the quotient polytope to be a cube.

Example 4.5. Let M be the 4-dimensional quasitoric manifold over a $2k$-gon with characteristic matrix of the form $\begin{pmatrix} 1 & 0 & 1 & 0 & \dots & 1 & 0 \\ 0 & 1 & 0 & 1 & \dots & 0 & 1 \end{pmatrix}$. Corollary 4.2 shows that the circle subgroup determined by the vector $\nu = (1,1)$ acts semifreely on M, but the quotient of M is not a 2-cube if $k > 2$, so M can not be homeomorphic to the product of spheres (it may be shown that M is the connected sum of $k-1$ copies of $S^2\times S^2$).

Surprisingly, quasitoric manifolds over polygons provide the only essential source of counterexamples, see Theorem 4.8 below for the precise statement.

Lemma 4.6. *A simple polytope P of dimension $n \geqslant 2$ all of whose 2-faces are 4-gons is combinatorially equivalent to a cube.*

Proof. We may assume by induction that all facets of P are cubes, and prove that P is a cube. The statement can be found in [14, Exercise 0.1], but we include the proof for the sake of completeness. We shall prove the dual statement about simplicial polytopes. The simplicial polytope dual to P is a crosspolytope; we call its boundary K (which is a sphere triangulation) a *cross-complex*. Recall that the *star* of a vertex v in a simplicial complex K is the subcomplex $\operatorname{st} v$ consisting of all simplices that together with v span a simplex, and *link* of v is the subcomplex $\operatorname{lk} v \subset \operatorname{st} v$ consisting of simplices not containing v. The duality between P and K extends to the duality between the facets of P (which are $(n-1)$-dimensional simple polytopes) and the links of vertices of K (which are triangulations of $(n-2)$-spheres). The dual statement follows from the next lemma (which is slightly more general, as we do not assume K to be the boundary of a simplicial polytope there). $\square$

Lemma 4.7. *Let K be a connected simplicial complex of dimension $k \geqslant 2$. If the link of each vertex of K is a crosscomplex of dimension $k-1$, then K is a crosscomplex.*

Proof. Let v be a vertex of K. By the assumption, $\operatorname{lk} v$ is a crosscomplex of dimension $k-1$, so every vertex $p \in \operatorname{lk} v$ has a unique vertex $q \in \operatorname{lk} v$ which is not joined to p by an edge in $\operatorname{lk} v$. Still, p and q may be joined by an edge in K, so we consider the two cases.

Case 1. Suppose that there is a pair of vertices p, q in $\operatorname{lk} v$ that are not joined by an edge in K. Let $\mathcal{R}$ be the set of other vertices of $\operatorname{lk} v$. The cardinality of $\mathcal{R}$ is $2(k-1)$. The link $\operatorname{lk} p$ is a crosspolytope, so it has $2k$ vertices, and contains v and all elements in $\mathcal{R}$. Since q is not joined to p by an edge in K, q is not in $\operatorname{lk} p$; so there is another vertex $p' \in \operatorname{lk} p,\ p' \notin v \cup \mathcal{R}$. Similarly, we have $q' \in \operatorname{lk} q,\ q' \notin v \cup \mathcal{R}$. Now take any $r \in \mathcal{R}$ and consider $\operatorname{lk} r$. Since $\operatorname{lk} v$ is a $(k-1)$-dimensional crosscomplex, r is joined to $2(k-1)$ vertices of $\operatorname{lk} v$ by edges in $\operatorname{lk} v$. We also know that r is joined to v, p' and q'. But since $\operatorname{lk} r$ is also a $(k-1)$-dimensional crosscomplex, r may be joined to only $2k$ vertices. Therefore $p' = q'$, which implies that K is a crosscomplex.

Case 2. Suppose that every pair of vertices in $\operatorname{lk} v$ is joined by an edge in K. This will lead us to a contradiction. Each vertex u in $\operatorname{lk} v$ is joined to v and all vertices in $\operatorname{lk} v$ except u itself. These are all vertices joined to u by edges, because the number of vertices in $\operatorname{lk} u$ is $2k$. This means that any pair of vertices in K is joined by an edge, and that K has exactly $2k+1$ vertices. The number of k-simplices meeting at each vertex is 2^k, and a k-simplex has $k+1$ vertices. So the total number of k-simplices in K is $2^k(2k+1)/(k+1)$.

Now we calculate the total number of k-simplices in K in a different way. Let σ be a k-simplex of K not containing v. Then σ contains a pair of vertices, say p and q, that are not joined by an edge in $\operatorname{lk} v$ (otherwise σ itself must be in $\operatorname{lk} v$, since $\operatorname{lk} v$ is a crosscomplex). Let L be the link of p in $\operatorname{lk} v$. Then L is a crosscomplex of dimension $k-2$, and it also coincides with the link of q in $\operatorname{lk} v$. We have that $\operatorname{lk} p$ is the join $L * \{v, q\}$, because both subcomplexes have the same vertex sets and both are crosscomplexes, and similarly $\operatorname{lk} q = L * \{v, p\}$. Since σ contains p and q, it follows that $\sigma = \tau * p * q$ for some $(k-2)$-simplex $\tau \in L$. Therefore, σ has at least two faces of dimension $(k-1)$ in $\operatorname{lk} v$, namely $\tau * p$ and $\tau * q$. Neither of these can be a face of another k-simplex not containing v, because every $(k-1)$-simplex of K is a face of exactly two k-simplices by the assumption, and $\tau * p$ is also a face of $\tau * p * v$ and $\tau * q$ is also a face of $\tau * q * v$. It follows that the number of k-simplices not containing v is no more than half of the number of $(k-1)$-simplices in $\operatorname{lk} v$. The number of k simplices containing

v equals the number of $(k-1)$-simplices in $\operatorname{lk} v$. The latter number is 2^k, so the total number of k-simplices in K is $\leqslant 2^{k-1}+2^k$, which is less than $2^k(2k+1)/(k+1)$ if $k \geqslant 2$. This contradiction finishes the proof. □

Remark. Another proof of Lemma 4.6 may be given by establishing a non-degenerate simplicial map from K to a crosscomplex. Such a map is a topological (non-ramified) covering of a sphere by a sphere, so it must be an isomorphism for $n \geqslant 3$. This approach was used in [16].

Theorem 4.8. *Assume that a quasitoric manifold M admits a semifreely acting subcircle with isolated fixed points, and every 2-face of the quotient polytope P is a 4-gon. Then M is $\mathbb{S}^1$-equivariantly homeomorphic to a product of 2-dimensional spheres.*

Proof. By Lemma 4.6, the orbit polytope is a cube. By Theorem 4.3, M is equivalent to a Bott tower. Finally, by Theorem 4.4 it is $\mathbb{S}^1$-homeomorphic to a product of spheres. □

We are also able to deduce the main result of Ilinskii [16]:

Corollary 4.9. *A toric manifold X with a circle subgroup acting semifreely and with isolated fixed points is diffeomorphic to a product of 2-spheres.*

Proof. By Theorem 4.4, it suffices to show that X is a Bott tower. To do this we apply Corollary 3.5, that is, we show that the fan corresponding to X is combinatorially equivalent to the fan over a crosspolytope. The semifree circle subgroup acting on X also acts semifreely and with isolated fixed points on every characteristic submanifold X_j of X. Using induction by dimension and Lemma 4.7 we reduce the statement to the complex 2-dimensional case, that is, we need to show that the quotient polytope of a complex 2-dimensional toric manifold with a semifree circle subgroup action and isolated fixed points is a 4-gon. (Note that a complex 2-dimensional toric manifold is always projective, so that we can work with polytopes instead of fans.) The following case-by-case analysis is just a reformulation of the argument from [16, §3]; we give it here for the sake of completeness.

Let Σ be the fan corresponding to our complex 2-dimensional toric manifold. The one dimensional cones of Σ correspond to the facets (or edges) of the quotient polygon P^2. We need to show that there are exactly 4 of them. The values of the characteristic function on the facets of P^2 are given by the primitive vectors generating the corresponding one-dimensional cones of Σ. Let ν be the vector generating the semifree circle subgroup. We may choose the initial vertex p of P^2 so that ν belongs to the 2-dimensional cone of Σ corresponding to p. Then we index the primitive vectors λ_i, $1 \leqslant i \leqslant m$, so that ν is in the cone generated by λ_1 and λ_2, and any two consecutive vectors span a two-dimensional cone (see Fig. 1). This provides us with a refined $2 \times m$ characteristic matrix $\mathit{\Lambda}$. We have $\lambda_1=(1,0)$ and $\lambda_2=(0,1)$, and applying the criterion from Corollary 4.2 to the first cone $\langle \lambda_1, \lambda_2 \rangle$ (i.e., to the initial vertex of the polygon), we obtain $\nu=(1,1)$.

Now consider the second cone. The non-singularity condition gives us $\det(\lambda_2,\lambda_3)=1$, whence $\lambda_3=(-1,*)$. Writing $\nu=k_1\lambda_2+k_2\lambda_3$ and applying Corollary 4.2 to the second cone $\langle \lambda_2, \lambda_3 \rangle$, we obtain

$$(1,1)=\pm(0,1)\pm(-1,*).$$



FIGURE 1.

Therefore, $\lambda_3 = (-1,0)$ or $\lambda_3 = (-1,-2)$. Similarly, considering the last cone $\langle \lambda_m, \lambda_1 \rangle$, we get $\lambda_m = (*,-1)$, and then, applying Corollary 4.2, $\lambda_m = (0,-1)$ or $\lambda_m = (-2,-1)$. The case $\lambda_3 = (-1,-2)$ and $\lambda_m = (-2,-1)$ is impossible as then the second and the last cones overlap.

Assume $\lambda_3 = (-1,0)$. Then a similar analysis applied to the third cone $\langle \lambda_3, \lambda_4 \rangle$ shows that $\lambda_4 = (0,-1)$ or $\lambda_4 = (-2,-1)$. Therefore, $\lambda_4 = \lambda_m$ (otherwise cones overlap).

Similarly, if $\lambda_m = (0,-1)$, we get $\lambda_{m-1} = (-1,0)$ or $\lambda_{m-1} = (-1,-2)$. Therefore, $\lambda_{m-1} = \lambda_3$.

In any case, we have $m = 4$ and P^2 is a 4-gon. This completes the proof. □

Note that the proof above leaves three possibilities for the vectors λ_3 and λ_4 of the corresponding 2-dimensional fan: $(-1,0)$ and $(0,-1)$, or $(-1,0)$ and $(-2,-1)$, or $(-1,-2)$ and $(0,-1)$, and the last two pairs are equivalent by an orientation reversing automorphism of T^2. The corresponding reduced submatrices are $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\begin{pmatrix} -1 & -2 \\ 0 & -1 \end{pmatrix}$. The first one corresponds to $\mathbb{C}P^1 \times \mathbb{C}P^1$, and the second to a non-trivial Bott tower (Hirzebruch surface) with $a_{12} = -2$. We finish this section by determining the class of matrices (2.2) corresponding to our specific class of Bott towers explicitly, for arbitrary dimension.

Theorem 4.10. *A Bott tower B^{2n} admits a semifree circle subgroup with isolated fixed points if and only if its matrix* (2.2) *satisfies the identity*

$$\frac{1}{2}(E - A) = C_1 C_2 \cdots C_n$$

where C_k is either a unit matrix or a unipotent upper triangular matrix with only one non-zero element $c_{i_k k} = 1$ in the k-th column off the diagonal, for $1 \leqslant k \leqslant n$.

Proof. First assume that B^{2n} admits a semifree circle subgroup with isolated fixed points. We have two sets of multiplicative generators for $H^*(B^{2n})$: the set $\{u_1, \ldots, u_n\}$ from Lemma 2.1, satisfying (2.1), and the set $\{x_1, \ldots, x_n\}$, satisfying $x_i^2 = 0$, $1 \leqslant i \leqslant n$. The reduced sets with $i \leqslant k$ can be considered as the corresponding sets of generators for the k-th stage B^{2k}. As it is clear from the proof of Theorem 4.4, we have $c_1(\xi_{k-1}) = -2c_{i_k k} x_{i_k}$ for some $i_k < k$, where $c_{i_k k} = 1$ or 0. From $u_k^2 + 2c_{i_k k}x_{i_k}u_k = 0$ we get $x_k = u_k + c_{i_k k}x_{i_k}$. In other words, the transition matrix C_k from the basis $x_1, \ldots, x_{k-1}, u_k, \ldots, u_n$ of $H^2(B^{2n})$ to $x_1, \ldots, x_k, u_{k+1}, \ldots, u_n$ may have only one non-zero entry off the diagonal, which is $c_{i_k k}$. The transition matrix from $u_1, \ldots, u_n$ to $x_1, \ldots, x_n$ is therefore the product $D = C_1 C_2 \cdots C_n$ (note that C_1 is the unit matrix as $x_1 = u_1$). Then $D = (d_{jk})$ is a unipotent upper triangular matrix consisting of zeros and units, we have $x_k = \sum_{j=1}^{n} d_{jk}u_j$, and

$$0 = x_k^2 = \bigl(u_k + \textstyle\sum_{j=1}^{k-1} d_{jk}u_j\bigr)^2 = u_k^2 + 2\textstyle\sum_{j=1}^{k-1} d_{jk}u_j u_k + \ldots, \quad k = 1, \ldots, n.$$

On the other hand, $0 = u_k^2 - \sum_{j=1}^{k-1} a_{jk}u_j u_k$ by (2.1). Comparing the coefficients of $u_j u_k$ for $1 \leqslant j \leqslant k-1$ in the last two equations and noting that these elements form a basis of the quotient space $H^4(B^{2k})/H^4(B^{2(k-1)})$, we obtain $2d_{jk} = -a_{jk}$ for $1 \leqslant j < k \leqslant n$. As both D and $-A$ are unipotent upper triangular matrices, this implies $2D = E - A$, as needed.

On the other hand, if A satisfies $E - A = 2C_1C_2 \cdots C_n$, then for the corresponding Bott tower we have $\xi_{k-1} = \pi_{i_k}^*(\gamma^{-2c_{i_k k}})$. Therefore we may choose a circle subgroup in such a way that ξ_{k-1} becomes $t\pi_{i_k}^*(\gamma^{-2c_{i_k k}})$ as an $\mathbb{S}^1$-bundle, for $1 < k \leqslant n$. This circle subgroup acts semifreely and with isolated fixed points by the same argument as in the proof of Theorem 4.4.

□

Example 4.11. It follows from Theorem 4.10 that if a Bott tower admits a semifree circle subgroup with isolated fixed points, then matrix (2.2) may have only 0 or -2 entries above the diagonal. However, the condition of Theorem 4.10 is stronger. For $n = 3$, the matrix

$$A = \begin{pmatrix} -1 & 0 & -2 \\ 0 & -1 & -2 \\ 0 & 0 & -1 \end{pmatrix}$$

does not occur for our class of Bott towers (since $(E - A)/2$ cannot be decomposed as $C_1C_2C_3$), while any other matrix with 0 or -2 above the diagonal occurs. For example, if

$$A = \begin{pmatrix} -1 & -2 & -2 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

then we have

$$\frac{1}{2}(E - A) = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

It is clear that not every Bott tower homeomorphic to a product of spheres admits a semifree subcircle action with isolated fixed points (the latter condition is stronger even for $n = 2$). We shall consider the class of Bott towers that are homeomorphic to a product of spheres in the next section.

5. TOPOLOGICAL CLASSIFICATION AND COHOMOLOGY

The following statement shows that Bott towers diffeomorphic to products of spheres can be detected by their cohomology rings, thereby providing a partial answer to Problem 2.3:

Theorem 5.1. *A Bott tower B^{2n} is diffeomorphic to $(\mathbb{C}P^1)^n$ if $H^*(B^{2n}) \cong H^*((\mathbb{C}P^1)^n)$ as rings.*

Proof. From Lemma 2.1 we get

$$H^*(B^{2n}) = H^*\big(B^{2n-2}\big)[u_n]/\big(u_n^2 - c_1(\xi_{n-1})u_n\big).$$

We may therefore write any element of $H^2(B^{2n})$ as $x + bu_n$ where $x \in H^2(B^{2n-2})$ and $b \in \mathbb{Z}$. Since

$$(x + bu_n)^2 = x^2 + 2bxu_n + b^2u_n^2 = x^2 + b(2x + bc_1(\xi_{n-1}))u_n,$$

the square of $x + bu_n$ with $b \neq 0$ is zero if and only if $x^2 = 0$ and $2x + bc_1(\xi_{n-1}) = 0$. This shows that the elements $x + bu_n$ with $b \neq 0$ whose squares are zero span a rank-one subgroup in $H^2(B^{2n})$.

By assumption, there is a basis $\{x_1, \ldots, x_n\}$ of $H^2(B^{2n})$ such that $x_i^2 = 0$ for all i. By the observation above, we may assume that $x_1, \ldots, x_{n-1}$ are in $H^2(B^{2n-2})$. Because $\{x_1, \ldots, x_n\}$ is a basis, x_n is not in $H^2(B^{2n-2})$ and we may assume $x_n = \sum_{i=1}^{n-1} b_ix_i + u_n$ with some $b_i \in \mathbb{Z}$. A product of the form $\prod_{i\in I} x_i$, where I is a subset of $\{1, \ldots, n\}$, belongs to $H^*(B^{2n-2})$ if and only if $n \notin I$. This shows that $H^*(B^{2n-2})$ is generated by $x_1, \ldots, x_{n-1}$ and has the same cohomology ring as $(\mathbb{C}P^1)^{n-1}$. Therefore, we may assume $B^{2n-2} \cong (\mathbb{C}P^1)^{n-1}$ by induction.

Writing $c_1(\xi_{n-1}) = \sum_{i=1}^{n-1} a_ix_i$, we see that

$$0 = x_n^2 = \big(u_n + \textstyle\sum_{i=1}^{n-1} b_ix_i\big)^2 = \sum_{i=1}^{n-1}(a_i + 2b_i)x_iu_n + \big(\sum_{i=1}^{n-1} b_ix_i\big)^2.$$

This may hold only if at most one a_i is non-zero because x_ix_j $(i < j)$ and x_iu_n form an additive basis of $H^4(B^{2n})$. Therefore, ξ_{n-1} is the pullback of γ^{-2b_i} over $\mathbb{C}P^1$ by a projection $B^{2n-2} = (\mathbb{C}P^1)^{n-1} \to \mathbb{C}P^1$. Since $P(\underline{\mathbb{C}} \oplus \gamma^{-2b_i})$ is a product bundle (see Example 2.2), so is $B^{2n} = P(\underline{\mathbb{C}} \oplus \xi_{n-1})$. □

We can now also effectively describe the class of matrices (2.2) corresponding to Bott towers which are diffeomorphic to a product of 2-spheres.

Theorem 5.2. *A Bott tower B^{2n} is diffeomorphic to $(\mathbb{C}P^1)^n$ if and only if the corresponding matrix (2.2) satisfies the identity*

$$\frac{1}{2}(E - A) = C_1C_2\cdots C_n$$

where each C_k is a unipotent upper triangular matrix which may have only one non-zero element in the k-th column off the diagonal, for $1 \leqslant k \leqslant n$.

Proof. We use the same argument as in the proof of Theorem 4.10. The only difference is that c_{i_kk} in $c_1(\xi_{k-1}) = -2c_{i_kk}x_{i_k}$ may now be an arbitrary integer. □

In the rest of this section we generalise the result of Theorem 5.1 to arbitrary quasitoric manifold, but only in the topological category (see Theorem 5.7).

We start by analysing the algebraic structure of the cohomology of quasitoric manifolds over a cube. Although it is possible to make this analysis over $\mathbb{Z}$, it is more convenient for our purposes to reduce the coefficients modulo 2. Let S be a graded algebra over $\mathbb{Z}/2$ generated by degree one elements $x_1, \dots, x_n$. We refer to S as a *Bott quadratic algebra* (simply *BQ-algebra*) of rank n if it satisfies the following two two properties:

(P1) $x_k^2 = \sum_{i<k} a_{ik} x_i x_k$ with $a_{ik} \in \mathbb{Z}/2$ for $1 \leqslant k \leqslant n$. (In particular, $x_1^2 = 0$.)

(P2) $\prod_{i=1}^n x_i \neq 0$.

If B^{2n} is a Bott tower, then (2.1) implies that $H^*(B^{2n}; \mathbb{Z}/2)$ is a *BQ*-algebra with the degree doubled, which explains our terminology. Our arguments below work for a wider class of algebras with (P1) replaced by the weaker property:

(P1') $x_k^2 = \sum_{i<j\leqslant k} a_{ijk} x_i x_j$ with $a_{ijk} \in \mathbb{Z}/2$ for $1 \leqslant k \leqslant n$.

Because of (P1) we can express any element of S as a linear combination of square-free monomials. We denote such a monomial $x_{i_1} \dots x_{i_s}$ by x_I, where $I = \{i_1, \dots, i_s\}$.

Lemma 5.3. *The elements x_I for all subsets $I \in \{1, \dots, n\}$ form an additive basis of S. In particular, $\dim S_q = \binom{n}{q}$ where S_q denotes the graded component of degree q.*

Proof. (P1) implies that the set $\{x_I\}$ additively generates S. We order monomials x_I using the inverse lexicographical order on subsets of $\{1, \dots, n\}$. Namely, if $I = \{i_1, \dots, i_s\}$ with $i_1 < \dots < i_s$ and similarly $J = \{j_1, \dots, j_s\}$, then we set $x_I < x_J$ if $i_k < j_k$ and $i_q = j_q$ for $k + 1 \leqslant q \leqslant s$.

Suppose that there is a non-trivial linear relation for x_I, and let x_J be the maximal monomial appearing in this relation. Then we may use the relation to replace the subfactor x_J in $\prod_{i=1}^n x_i$, and then use (P1) repeatedly whenever x_k^2 appears, until we end up at zero. This contradicts (P2). Therefore, there are no non-trivial linear relations among x_I's. □

Lemma 5.4. *Suppose we have a surjective graded homomorphism f from S to a graded algebra S' over $\mathbb{Z}/2$ satisfying $S'_{n-1} \neq 0$. Then the dimension of the kernel of $f\colon S_1 \to S'_1$ is at most one. Moreover, if the dimension of the kernel is exactly one, then S' is a BQ-algebra of rank $n - 1$.*

Proof. We denote $f(x_i)$ by $\bar{x}_i$. Then (P1) holds for $\bar{x}_1, \dots, \bar{x}_n$. Suppose the dimension of the kernel is at least two. Then there exist $p > q \geqslant 1$ such that

$$\bar{x}_p = \sum_{i<p} b_i \bar{x}_i, \qquad \bar{x}_q = \sum_{j<q} c_j \bar{x}_j \tag{5.1}$$

where $b_i, c_j \in \mathbb{Z}/2$. By Lemma 5.3, S_{n-1} is generated by the elements x_I with $|I| = n-1$. We shall show that $\bar{x}_I = 0$ for any such I, which contradicts the assumption $S'_{n-1} \neq 0$.

First assume $q \geqslant 2$. Since $|I| = n - 1$, I contains p or q. We replace $\bar{x}_p$ and $\bar{x}_q$ in $\bar{x}_I$ by (5.1) and use (P1) repeatedly whenever $\bar{x}_k^2$ appears. We end up at zero.

If $q = 1$, i.e., $\bar{x}_1 = 0$, then it suffices to see that $\bar{x}_I = 0$ for $I = \{2, 3, \dots, n\}$. We replace $\bar{x}_p$ in $\bar{x}_I$ by (5.1) and use (P1) repeatedly whenever $\bar{x}_k^2$ appears for $k \geqslant 2$. Then each term in the final expression contains a factor $\bar{x}_1$, which is zero.

Now we prove the second statement of the lemma. By the assumption, the elements $\bar{x}_i$ satisfy a non-trivial linear relation. Let $\bar{x}_j$ be the maximal element appearing in this relation. We can eliminate $\bar{x}_j$ in S' using this linear relation and (P1). Then (P1) holds for $\bar{x}_i$'s with $i \neq j$. Therefore, S'_{n-1} is generated by $\prod_{i\neq j} \bar{x}_i$. This element is non-zero because $S'_{n-1} \neq 0$. This proves that S' is a BQ-algebra of rank $n-1$. □

Theorem 5.5. *Let M be a quasitoric manifold with the quotient polytope P. Then $H^{2*}(M;\mathbb{Z}/2)$ is a BQ-algebra of rank n if and only if P is an n-cube.*

Proof. Assume that P is an n-cube. Since every principal minor of $\Lambda_\star$ is 1 modulo 2, we deduce that $\Lambda_\star$ is conjugate to a unipotent upper triangular matrix by the same argument as in the proof of Lemma 3.3. (Matrix (3.7) in which all b_i are non-zero modulo 2 for $1 \leqslant i \leqslant n$ cannot occur because its determinant is zero modulo 2.) Then it follows from from (3.4) that $H^{2*}(M;\mathbb{Z}/2)$ is a BQ-algebra of rank n.

Now assume that $H^{2*}(M;\mathbb{Z}/2)$ is a BQ-algebra. Let $b_r(M)$ denote the r-th Betti number of M, and let $f_s(P)$ denote the number of faces of P of codimension $s+1$. Then

$$b_2(M) = f_0(P) - n, \qquad b_4(M) = f_1(P) - (n-1)f_0(P) + \binom{n}{n-2}$$

(see [7, Th. 3.1]), and we obtain from Lemma 5.3 that

$$f_0(P) = 2n, \qquad f_1(P) = 2n(n-1). \tag{5.2}$$

For every characteristic submanifold M_i the restriction map $H^*(M;\mathbb{Z}/2) \to H^*(M_i;\mathbb{Z}/2)$ is surjective [12, Lemma 2.3]. It follows from Lemmas 5.3 and 5.4 that $b_2(M_i) \geqslant b_2(M) - 1 = n-1$. Therefore,

$$f_0(F_i) = (n-1) + b_2(M_i) \geqslant 2(n-1), \tag{5.3}$$

where F_i denotes the facet corresponding to M_i, and

$$f_1(P) = \frac{1}{2}\sum_{i=1}^{2n} f_0(F_i) \geqslant 2n(n-1).$$

Comparing this with (5.2) we obtain that the equality holds in (5.3) for every i, that is, $b_2(M_i) = n-1$. This implies that the kernel of $H^2(M;\mathbb{Z}/2) \to H^2(M_i;\mathbb{Z}/2)$ is one-dimensional, so $H^{2*}(M_i;\mathbb{Z}/2)$ is a BQ-algebra of rank $n-1$ by Lemma 5.4. This enables us to use induction on n.

When $n = 2$, equations (5.2) imply that P is combinatorially a square. Suppose that the theorem holds for $n-1$ with $n \geqslant 3$. Since $H^{2*}(M_i)$ is a BQ-algebra of rank $n-1$, every facet of P is an $(n-1)$-cube; in particular, every 2-face of P is a square. Then P is an n-cube by virtue of Lemma 4.6. □

Lemma 5.6. *Let M be a quasitoric manifold over an n-cube. If $H^*(M;\mathbb{Q}) \cong H^*((\mathbb{C}P^1)^n;\mathbb{Q})$ as rings, then M is equivalent to a Bott tower.*

Proof. By the assumption, there are elements $y_1, \dots, y_n$ in $H^2(M;\mathbb{Q})$ generating $H^*(M;\mathbb{Q})$ and satisfying $y_i^2 = 0$ for $1 \leqslant i \leqslant n$. Let $M_i \subset M$ be a characteristic submanifold, and denote the restriction of y_k to $H^2(M_i;\mathbb{Q})$

by $\bar{y}_k$. Then $\bar{y}_1, \dots, \bar{y}_n$ generate $H^*(M_i;\mathbb{Q})$ as a ring because $H^*(M;\mathbb{Q}) \to H^*(M_i;\mathbb{Q})$ is surjective. Since $b_2(M_i) = n-1$, there is a non-trivial linear relation for the elements $\bar{y}_k$. Using this linear relation, we can eliminate one generator, say $\bar{y}_n$, and get a surjective map $\mathbb{Q}[\bar{y}_1, \dots, \bar{y}_{n-1}]/(\bar{y}_1^2, \dots, \bar{y}_{n-1}^2) \to H^*(M_i;\mathbb{Q})$. Since the dimensions of components of degree $2q$ in both rings equal $\binom{n-1}{q}$, the surjective map is an isomorphism. Therefore, $H^*(M_i;\mathbb{Q}) \cong H^*((\mathbb{C}P^1)^{n-1};\mathbb{Q})$, so we may use an induction argument and assume that every M_i is a Bott tower.

Let $\Lambda_\star$ be the reduced submatrix of M. It follows from Lemma 3.3 that $-\Lambda_\star$ is conjugate to a unipotent upper triangular matrix or to matrix (3.7) with non-zero entries b_i for $1 \leqslant i \leqslant n$. It suffices to exclude the latter. Suppose that $-\Lambda_\star$ is given by (3.7). Then $\det(-\Lambda_\star) = -1$, that is,

$$\prod_{i=1}^{n} b_i = (-1)^n 2. \tag{5.4}$$

Using (3.4), we obtain

$$H^*(M) = \mathbb{Z}[x_1, \dots, x_n]/\big(x_1(x_1+b_1x_2), x_2(x_2+b_2x_3), \dots, x_n(b_nx_1+x_n)\big),$$

where we set $x_i = v_{i+n}$ for $1 \leqslant i \leqslant n$. By the assumption, there is a non-zero element $x \in H^2(M,\mathbb{Q})$ whose square is zero. Write $x = \sum_{i=1}^n a_ix_i$ for some $a_i \in \mathbb{Q}$, then

$$\begin{aligned} 0 = \Big(\sum_{i=1}^n a_ix_i\Big)^2 &= \sum_{i=1}^n a_i^2x_i^2 + 2\sum_{i<j} a_ia_jx_ix_j \\ &= -a_1^2b_1x_1x_2 - a_2^2b_2x_2x_3 - \ldots - a_n^2b_nx_nx_1 + 2\sum_{i<j} a_ia_jx_ix_j, \end{aligned}$$

which implies that

$$a_1^2b_1 = 2a_1a_2, \quad a_2^2b_2 = 2a_2a_3, \quad \dots, \quad a_n^2b_n = 2a_na_1. \tag{5.5}$$

Suppose $a_i \neq 0$ for every i. Multiplying the identities above, we obtain $\prod_{i=1}^n b_i = 2^n$, which contradicts (5.4). Therefore $a_i = 0$ for some i, but this together with (5.5) implies that $a_i = 0$ for every i. This contradicts the assumption that $x = \sum_{i=1}^n a_ix_i \neq 0$. Therefore, (3.7) cannot occur as a reduced characteristic matrix, and M is a Bott tower. □

We are now ready to prove our final promised result.

Theorem 5.7. *A quasitoric manifold M is homeomorphic to $(\mathbb{C}P^1)^n$ if and only if $H^*(M) \cong H^*((\mathbb{C}P^1)^n)$ as rings.*

Proof. Since $H^*((\mathbb{C}P^1)^n;\mathbb{Z}/2)$ is a BQ-algebra of rank n, the quotient polytope of M is an n-cube by Theorem 5.5. Then M is a Bott tower by Lemma 5.6. Finally, M is homeomorphic to $(\mathbb{C}P^1)^n$ by Theorem 5.1. □

We finish by proposing the following quasitoric analogue of Problem 2.3.

Problem 5.8. *Does an isomorphism of rings $H^*(M_1) \cong H^*(M_2)$ imply a homeomorphism of quasitoric manifolds M_1 and M_2?*

References

[1] Raoul Bott and Hans Samelson. *Application of the theory of Morse to symmetric spaces.* Amer. J. of Math. **80** (1958), 964–1029.

[2] Victor M. Buchstaber and Taras E. Panov. *Torus Actions and Their Applications in Topology and Combinatorics.* University Lecture Series, vol. **24**, Amer. Math. Soc., Providence, R.I., 2002.

[3] Victor M. Buchstaber, Taras E. Panov and Nigel Ray. *Spaces of polytopes and cobordism of quasitoric manifolds.* Moscow Math. J. **7** (2007), no. 2; arXiv:math.AT/0609346.

[4] Victor M. Buchstaber and Nigel Ray. *Tangential structures on toric manifolds, and connected sums of polytopes.* Internat. Math. Res. Notices **4** (2001), 193–219; arXiv:math.AT/0010025.

[5] Suyoung Choi, Mikiya Masuda and Dong Youp Suh. *Quasi-toric manifolds and small covers over a product of simplices.* In preparation, 2007.

[6] Yusuf Civan and Nigel Ray. *Homotopy decompositions and K-theory of Bott towers.* K-Theory **34** (2005), no. 1, 1–33.

[7] Michael W. Davis and Tadeusz Januszkiewicz. *Convex polytopes, Coxeter orbifolds and torus actions.* Duke Math. J. **62** (1991), no. 2, 417–451.

[8] Michael Grossberg and Yael Karshon. *Bott towers, complete integrability, and the extended character of representations.* Duke Math. J. **76** (1994), 23–58.

[9] Akio Hattori. *Symplectic manifolds with semi-free Hamiltonian S^1-action.* Tokyo J. Math. **15** (1992), no. 2, 281–296.

[10] Akio Hattori and Tomoyoshi Yoshida. *Lifting compact group actions in fiber bundles.* Japan. J. Math. **2** (1976), 13–25.

[11] Mikiya Masuda. *Unitary toric manifolds, multi-fans and equivariant index.* Tohoku Math. J. (2) **51** (1999), no. 2, 237–265.

[12] Mikiya Masuda and Taras Panov. *On the cohomology of torus manifolds.* Osaka J. Math. **43** (2006), 711–746; arXiv:math.AT/0306100.

[13] Susan Tolman and Jonathan Weitsman. *On semifree symplectic circle actions with isolated fixed points.* Topology **39** (2000), no. 2, 299–309.

[14] Günter M. Ziegler. *Lectures on polytopes.* Graduate Texts in Mathematics, vol. **152**, Springer-Verlag, New York, 1995.

[15] Н. Э. Добринская. *Проблема классификации квазиторических многообразий над заданным простым многогранником.* Функц. анализ и его прил. **35** (2001), вып. 2, 3–11 (in Russian). English translation: Natalia Dobrinskaya. *Classification problem for quasitoric manifolds over a given simple polytope.* Funct. Anal. Appl. **35** (2001), no. 2, 83–89.

[16] Д. Г. Ильинский. *О почти свободном действии одномерного тора на торическом многообразии.* Мат. Сборник **197** (2006), вып. 5, 51–74 (in Russian). English translation: Dmitri Ilinskii. *On the almost free action of one-dimensional torus on a toric variety.* Sbornik Math. **197** (2006), no. 5, 681–703.

[17] Т. Е. Панов. *Роды Хирцебруха многообразий с действием тора.* Изв. РАН, сер. матем **65** (2001), вып. 3, 123–138 (in Russian). English translation: Taras E. Panov. *Hirzebruch genera of manifolds with torus action.* Izvestiya. Math., **65** (2001), no. 3, 543–556; arXiv:math.AT/9910083.

Department of Mathematics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan

Department of Geometry and Topology, Faculty of Mathematics and Mechanics, Moscow State University, Leninskiye Gory, Moscow 119992, Russia

E-mail address: `masuda@sci.osaka-cu.ac.jp`

E-mail address: `tpanov@mech.math.msu.su`